
Ashton & Parsons' INFANTS' POWDERS
 tincture of matricaria

THR Holder: Alliance Pharmaceuticals Limited, SM15 2BB, UK.
 Manufacturer: Pharmapac (UK) Limited, CH41 7EL, UK.
 Ashton & Parsons, Alliance and associated devices are registered trademarks of Alliance Pharmaceuticals Limited

© Alliance Pharmaceuticals Limited 2017
 ALLIANCE
 Text revised February 2017
 THR 1683/0076

20 SACHET PACK
Ashton & Parsons' INFANTS' POWDERS
 tincture of matricaria
For teething pain relief
 For oral use

A traditional herbal medicinal product used for the relief of teething pain and symptoms associated with teething such as sore and tender gums, flushed cheeks and dribbling based on traditional use only



For teething pain relief
 For oral use

A traditional herbal medicinal product used for the relief of teething pain and symptoms associated with teething such as sore and tender gums, flushed cheeks and dribbling based on traditional use only

20 SACHET PACK
Ashton & Parsons' INFANTS' POWDERS
 tincture of matricaria

What these powders contain
 Each sachet contains powder impregnated with 0.002ml of the active ingredient tincture of matricaria (*Matricaria recutita* L.) flower. Also contains lactose. Extraction solvent: Ethanol 70% v/v

How to use these powders
 Not recommended for use in babies under 3 months. For use in teething infants only.

New instruction for use
 Ensure that the infant is held in a sitting or upright position. Put the powder slowly on the front of the infant's tongue, a little at a time, allowing the powder to fully dissolve.

Age	Recommended dose	Additional dosing
Infants aged 3-6 months	Use a teaspoon to give the powder. Half the contents of one sachet in the morning and the other half in the evening. The sachet should be divided in half by emptying half the contents onto the teaspoon and retaining the other half in the sachet.	If the infant is very restless, the dose may be repeated every 1, 2 or 3 hours until improvement occurs. Do not give more than 6 doses in 24 hours.
Infants over 6 months	One sachet in the morning and one sachet in the evening.	

Do not give more than the label tells you to. If symptoms persist for more than 7 days, speak to your doctor, if you give your child more than you should, it may cause diarrhoea. If the infant experiences any side effects, stop using this medicine and speak to your doctor or pharmacist.

Possible side effects
 Like all medicines, this medicine can cause side effects, although not everybody gets them.
Possible side effects: allergic reaction which may include rash, tongue sores.

Reporting of side effects
 If your child gets any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the internet at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.




Before using these powders
Do not use if your child


- is allergic to tincture of matricaria, lactose or to plants of the Asteraceae/Compositae family such as daisies, marigolds or antirrhines
- has an intolerance to some sugars, unless your doctor tells you to (this medicine contains lactose).

Warning: This medicine may cause an allergic reaction or make existing conditions worse in children who suffer from allergic type conditions (e.g. asthma).


How to store these powders
 Keep this medicine out of the sight and reach of children.
 Store this medicine below 25°C.
 Use by the expiry date on the sachet/carton (EXP).
 Do not use this medicine if before first use you find the end seals are broken.

5 036631 001685 >



BRaille CONVERSION - English Alphabet



ASHTON
AND
PARSONS

BRaille CONVERSION - English Alphabet



INFANTS
POWDERS



IC0180

How to store these powders

Keep this medicine out of the sight and reach of children.
 Store this medicine below 25°C.
 Use by the expiry date on the sachet/carton (EXP).
 Do not use this medicine if before first use you find the end seals are broken.


Ashton & Parsons®
INFANTS' POWDERS
 tincture of matricaria
For teething pain relief
 For oral use

A traditional herbal medicinal product used for the relief of teething pain and symptoms associated with teething such as sore and flushed cheeks and dribbling based on traditional use only



Ashton & Parsons' INFANTS' POWDERS
 tincture of matricaria

THR Holder: Alliance Pharmaceuticals Limited, SN15 2BB, UK.
 Manufacturer: Pharnapac (UK) Limited, CH41 7EL, UK.
 Trademarks of Alliance Pharmaceuticals Limited
 © Alliance Pharmaceuticals Limited 2017

THR 1683/0076 **ALLIANCE** Text revised February 2017

30 SACHET PACK
Ashton & Parsons' INFANTS' POWDERS
 tincture of matricaria
For teething pain relief
 For oral use



What these powders contain
 Each sachet contains powder impregnated with 0.002ml of the active ingredient tincture of matricaria (*Matricaria recutita* L.) flower. Also contains lactose. Extraction solvent: Ethanol 70% v/v

How to use these powders
 Not recommended for use in babies under 3 months. For use in teething infants only.

New instruction for use
 Ensure that the infant is held in a sitting or upright position. Put the powder slowly on the front of the infant's tongue, a little at a time, allowing the powder to fully dissolve.

Age	Recommended dose	Additional dosing
Infants aged 3-6 months	Use a teaspoon to give the powder. Half the contents of one sachet in the morning and the other half in the evening. The sachet should be divided in half by emptying half the contents onto the teaspoon and retaining the other half in the sachet.	If the infant is very restless, the dose may be repeated every 1, 2, or 3 hours until improvement occurs. Do not give more than 6 doses in 24 hours.
Infants over 6 months	One sachet in the morning and one sachet in the evening.	

Do not give more than the label tells you to. If symptoms persist for more than 7 days, speak to your doctor. **If you give your child more than you should, it may cause diarrhoea.** If the infant experiences any side effects, stop using this medicine and speak to your doctor or pharmacist.


Possible side effects
 Like all medicines, this medicine can cause side effects, although not everybody gets them.
Possible side effects: allergic reaction which may include rash, tongue sores.

Reporting of side effects
 If your child gets any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the internet at www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.


Before using these powders
Do not use if your child

- is allergic to tincture of matricaria, lactose or to plants of the Asteraceae/Compositae family such as daisies, margoldis or anemones
- has an intolerance to some sugars, unless your doctor tells you to (this medicine contains lactose).

Warning: This medicine may cause an allergic reaction or make existing conditions worse in children who suffer from allergic type conditions (e.g. asthma).



BRAILLE CONVERSION - English Alphabet



ASHTON
AND
PARSONS

BRAILLE CONVERSION - English Alphabet



INFANTS
POWDERS

