

## PACKAGE LEAFLET: INFORMATION FOR THE USER

### Epilim® 200mg and 500mg Gastro-resistant Tablets

sodium valproate

**Is this leaflet hard to see or read? Phone 0845 372 7101 for help.**

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

#### WARNING

Valproate can cause birth defects and problems with early development of the child if it is taken during pregnancy. If you are a female of childbearing age you should use an effective method of contraception throughout your treatment.

Your doctor will discuss this with you but you should also follow the advice in section 2 of this leaflet. Tell your doctor at once if you become pregnant or think you might be pregnant.

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, please ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them even if their symptoms are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet

1. What Epilim is and what it is used for
2. What you need to know before you take Epilim
3. How to take Epilim
4. Possible side effects
5. How to store Epilim
6. Contents of the pack and other information

### 1. What Epilim is and what it is used for

#### What Epilim is

The name of your medicine is Epilim 200mg or 500mg Gastro-resistant Tablets (called Epilim in this leaflet). Epilim 200mg or 500mg Gastro-resistant Tablets are “enteric coated” this means that the tablets have a protective coating that allows them to reach the intestines (gut) without being dissolved in the stomach first. This helps stop them from causing a stomach upset.

#### What Epilim contains

Epilim contains sodium valproate. It belongs to a group of medicines called anti-convulsants or anti-epileptic agents. It works by helping to calm the brain down.

#### What Epilim is used for

Epilim is used to treat epilepsy (fits) in adults and children.

## **2. What you need to know before you take Epilim**

### **Do not take Epilim and tell your doctor if:**

- X You are allergic (hypersensitive) to sodium valproate or any of the other ingredients of Epilim (see Section 6: Contents of the pack and other information).  
Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue.
- X You have liver problems or you or your family have a history of liver problems.
- X You have a rare illness called porphyria.
- X You have a known metabolic disorder, i.e. a urea cycle disorder.
- X You have a genetic problem caused by a mitochondrial disorder (e.g. Alpers-Huttenlocher syndrome).

Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Epilim.

### **Warnings and precautions**

A small number of people being treated with anti-epileptics such as sodium valproate have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

As with other anti-epileptic drugs, convulsions may become worse or happen more frequently whilst taking this medicine. If this happens contact your doctor immediately.

### **Talk to your doctor or pharmacist before taking Epilim if:**

- You have diabetes. This medicine may affect the results of urine tests.
- You have a carnitine palmitoyltransferase type II deficiency.
- You have kidney problems. Your doctor may give you a lower dose.
- You have fits (epilepsy), brain disease or a metabolic condition affecting your brain.
- You have a 'urea cycle disorder' where too much ammonia builds up in the body.
- You have an illness called 'systemic lupus erythematosus (SLE)' – a disease of the immune system which affects skin, bones, joints and internal organs.
- You know that there is a genetic problem caused by a mitochondrial disorder in your family.

If you are not sure if any of the above apply to you, talk to your doctor or pharmacist before taking Epilim.

### **Weight gain**

Taking Epilim may make you put on weight. Talk to your doctor about how this will affect you.

### **Blood tests**

Your doctor may wish to do blood tests before you start taking Epilim and during your treatment.

### **Other medicines and Epilim**

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines you buy without a prescription, including herbal medicines. This is because Epilim can affect the way some other medicines work. Also some medicines can affect the way Epilim works.

**The following medicines can increase the chance of you getting side effects, when taken with Epilim:**

- Some medicines used for pain and inflammation (salicylates) such as aspirin.
- Some other medicines used to treat fits (epilepsy) – see page 2, section 3, ‘Patients taking other medicines for fits’. This includes medicines such as phenobarbital, primidone, phenytoin, carbamazepine, rufinamide, topiramate, acetazolamide, lamotrigine and felbamate.

**Epilim may increase the effect of the following medicines:**

- Medicines used for thinning the blood (such as warfarin).
- Zidovudine – used for HIV infection.
- Temozolomide used to treat cancer.
- Medicines for depression.
- Monoamine oxidase inhibitors (MAOI) such as moclobemide, selegiline, linezolid.
- Medicines used to calm emotional and mental health problems (including schizophrenia, bipolar disorder and depression) such as quetiapine, diazepam and olanzapine.
- Nimodipine.
- Propofol – used for anaesthesia.

**The following medicines can affect the way Epilim works:**

- Some medicines used for the prevention and treatment of malaria such as mefloquine and chloroquine.
- Cimetidine – used for stomach ulcers.
- Protease inhibitors such as lopinavir and ritonavir – used for HIV infection and AIDS.
- Carbapenem agents (antibiotics used to treat bacterial infections) such as imipenem, meropenem, rifampicin and erythromycin. The combination of Epilim and carbapenems should be avoided because it may decrease the effect of your medicine.
- Cholestyramine used to lower blood fat (cholesterol) levels.

**Taking Epilim with food and drink**

Alcohol intake is not recommended during treatment.

**Pregnancy and breast-feeding**

**Important advice for women**

- Valproate can be harmful to unborn children when taken by a woman during pregnancy.
- Valproate carries a risk if taken during pregnancy. The higher the dose, the higher the risks but all doses carry a risk.
- It can cause serious birth defects and can affect the way in which the child develops as it grows. Birth defects which have been reported include *spina bifida* (where the bones of the spine are not properly developed); facial and skull malformations; heart, kidney, urinary tract and sexual organ malformations; limb defects.
- If you take valproate during pregnancy you have a higher risk than other women of having a child with birth defects that require medical treatment. Because valproate has been used for many years we know that in women who take valproate around 10 babies in every 100 will have birth defects. This compares to 2-3 babies in every 100 born to women who don't have epilepsy.
- It is estimated that up to 30-40% of preschool children whose mothers took valproate during pregnancy may have problems with early childhood development. Children affected can be slow to walk and talk, intellectually less able than other children, and have difficulty with language and memory.

- Autistic spectrum disorders are more often diagnosed in children exposed to valproate and there is some evidence children may be more likely to develop symptoms of Attention Deficit Hyperactivity Disorder (ADHD).
- If you are a woman capable of becoming pregnant your doctor should only prescribe valproate for you if nothing else works for you.
- Before prescribing this medicine to you, your doctor will have explained what might happen to your baby if you become pregnant whilst taking valproate. If you decide later you want to have a child you should not stop taking your medicine until you have discussed this with your doctor and agreed a plan for switching you onto another product if this is possible.
- Ask your doctor about taking folic acid when trying for a baby. Folic acid can lower the general risk of *spina bifida* and early miscarriage that exists with all pregnancies. However, it is unlikely that it will reduce the risk of birth defects associated with valproate use.

### **FIRST PRESCRIPTION**

If this is the first time you have been prescribed valproate your doctor will have explained the risks to an unborn child if you become pregnant. Once you are of childbearing age, you will need to make sure you use an effective method of contraception throughout your treatment. Talk to your doctor or family planning clinic if you need advice on contraception.

#### **Key messages:**

- Make sure you are using an effective method of contraception.
- Tell your doctor at once if you are pregnant or think you might be pregnant.

### **CONTINUING TREATMENT AND NOT TRYING FOR A BABY**

If you are continuing treatment with valproate but you don't plan to have a baby make sure you are using an effective method of contraception. Talk to your doctor or family planning clinic if you need advice on contraception.

#### **Key messages:**

- Make sure you are using an effective method of contraception.
- Tell your doctor at once if you are pregnant or think you might be pregnant.

### **CONTINUING TREATMENT AND CONSIDERING TRYING FOR A BABY**

If you are continuing treatment with valproate and you are now thinking of trying for a baby you must not stop taking either your valproate or your contraceptive medicine until you have discussed this with your prescriber. You should talk to your doctor well before you become pregnant so that you can put several actions in place so that your pregnancy goes as smoothly as possible and any risks to you and your unborn child are reduced as much as possible.

Your doctor may decide to change the dose of valproate or switch you to another medicine before you start trying for a baby.

If you do become pregnant you will be monitored very closely both for the management of your underlying condition and to check how your unborn child is developing.

Ask your doctor about taking folic acid when trying for a baby. Folic acid can lower the general risk of *spina bifida* and early miscarriage that exists with all pregnancies. However, it is unlikely that it will reduce the risk of birth defects associated with valproate use.

#### **Key messages:**

- Do not stop using your contraception before you have talked to your doctor and worked together on a plan to ensure your epilepsy is controlled and the risks to your baby are reduced.
- Tell your doctor at once when you know or think you might be pregnant.

### **UNPLANNED PREGNANCY WHILST CONTINUING TREATMENT**

Babies born to mothers who have been on valproate are at serious risk of birth defects and problems with development which can be seriously debilitating. If you are taking valproate and you think you are pregnant or might be pregnant contact your doctor at once. Do not stop taking your medicine until your doctor tells you to.

Ask your doctor about taking folic acid. Folic acid can lower the general risk of *spina bifida* and early miscarriage that exists with all pregnancies. However, it is unlikely that it will reduce the risk of birth defects associated with valproate use.

#### **Key messages:**

- Tell your doctor at once if you know you are pregnant or think you might be pregnant.
- Do not stop taking valproate unless your doctor tells you to.

**Make sure you read the patient booklet and sign the Acknowledgement of Risk form which should be given to you and discussed with you by your doctor or pharmacist.**

#### **Breast-feeding**

Very little Epilim gets into the breast milk. However, talk to your doctor about whether you should breast-feed your baby. Ask your doctor or pharmacist for advice before taking any medicine.

#### **Driving and using machines**

You may feel sleepy when taking Epilim. If this happens to you, do not drive or use any tools or machines. Taking other medicines used to treat fits or calm emotional and mental health problems may increase sleepiness.

### **3. How to take Epilim**

Always take Epilim exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Epilim treatment must be started and supervised by a doctor specialised in the treatment of epilepsy.

#### **Taking this medicine**

- Your doctor will decide how much Epilim to give you or your child depending on your or your child's body weight.
- Take this medicine by mouth.
- **Do not** crush or chew the tablets.
- If you feel the effect of your medicine is too weak or too strong, do not change the dose yourself but ask your doctor.

#### **How to take this medicine**

- The dose is normally split and given half in the morning and half in the evening.

#### **How much to take**

##### **Adults (including the elderly)**

- The starting dose is 600mg daily. Your doctor should gradually increase this dose by 200mg every 3 days depending on your condition.
- The usual dose is between 1000mg and 2000mg (20-30mg per kilogram of body weight) each day.
- This may be increased to 2500mg each day depending on your illness.

**Children over 20 kilograms**

- The starting dose should be 400mg daily. Your doctor should increase this dose depending on your child's illness.
- The usual dose is then between 20mg and 30mg for each kilogram of body weight each day.
- This may be further increased to 35mg for each kilogram of body weight each day depending on your child's illness.

**Children under 20 kilograms**

- The usual dose is 20mg for each kilogram of body weight each day.
- Depending on the child's condition your child's doctor may decide to increase this dose.

**Patients with kidney problems**

Your doctor may decide to adjust your or your child's dose.

**Patients taking other medicines for fits (epilepsy):**

- You or your child may be taking other medicines for epilepsy at the same time as Epilim. If so, your doctor should gradually initiate treatment depending on your or your child's condition.
- Your doctor may increase the dose of Epilim by 5 to 10mg for each kilogram of body weight each day depending on which other medicines you are taking.

**If you take more Epilim than you should**

If you take more Epilim than you should, tell a doctor or go to a hospital casualty department straight away. Take the medicine pack with you. This is so the doctor knows what you have taken.

The following effects may happen: feeling sick or being sick, pupils of the eye become smaller, dizziness, loss of consciousness, weak muscles and poor reflexes, breathing problems, headaches, fits (seizures), confusion, memory loss and unusual or inappropriate behaviour.

**If you forget to take Epilim**

If you forget to take a dose, take it as soon as you remember. However, if it is nearly time for the next dose, skip the missed dose. Do not take a double dose to make up for a forgotten dose.

**If you stop taking Epilim**

Keep taking until your doctor tells you to stop. Do not stop taking Epilim just because you feel better. If you stop your fits may come back.

**Tests**

Make sure you or your child keep your regular appointments for a check-up. They are very important as your or your child's dose may need to be changed. Epilim can change the levels of liver enzymes shown up in blood tests. This can mean that your or your child's liver is not working properly. If you or your child go into hospital or visit another doctor or a dentist, tell them you are taking Epilim.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, Epilim can cause side effects, although not everybody gets them.

**Tell your doctor straight away if you notice any of the following serious side effects – you may need urgent medical treatment:**

- You have an **allergic reaction**. The signs may include: a rash, joint pain, fever (systemic lupus erythematosus), swallowing or breathing problems, swelling of your lips, face, throat or tongue. Hands, feet or genitals may also be affected. More severe allergic reactions can lead to lymph node enlargement and possible impairment of other organs.
- Liver problems and problems of the pancreas may show as a sudden illness which may happen in the first six months of treatment. This happens in a very small number of people taking Epilim. It includes feeling and being sick many times, being very tired, sleepy and weak, stomach pain including very bad upper stomach pain, jaundice (yellowing of the skin or whites of the eyes), loss of appetite, swelling (especially of the legs and feet but may include other parts of the body), worsening of your fits or a general feeling of being unwell. Your doctor may tell you to stop taking Epilim immediately if you have these symptoms.
- You have a skin rash or skin lesions with a pink/red ring and a pale centre which may be itchy, scaly or filled with fluid. The rash may appear especially on the palms or soles of your feet. These could be signs of a serious allergy to the medicine called 'erythema multiforme'.
- Blistering or bleeding of the skin around the lips, eyes, mouth, nose and genitals. Also flu-like symptoms and fever. This may be something called 'Stevens-Johnson syndrome'.
- Severe blistering rash where layers of the skin may peel off to leave large areas of raw exposed skin over the body. Also a feeling of being generally unwell, fever, chills and aching muscles. This may be something called 'Toxic epidermal necrolysis'.
- Bruising more easily and getting more infections than usual. This could be a blood problem called 'thrombocytopenia'. It can also be due to a fall in the number of white blood cells, bone marrow depression or another condition that affects red blood cells, white blood cells and platelets (pancytopenia) or how the blood clots.
- Blood clotting problems (bleeding for longer than normal), bruising or bleeding for no reason.
- Changes in mood, loss of memory, lack of concentration and deep loss of consciousness (coma).
- Underactive thyroid gland, which may cause tiredness or weight gain (hypothyroidism).
- Breathing difficulty and pain due to inflammation of the lungs (pleural effusion).

**Tell your doctor as soon as possible if you have any of the following side effects:**

- Changes in behaviour including being very alert, and sometimes also aggressive, hyperactive and unusual or inappropriate behaviour. This is more likely if other medicine to treat fits such as phenobarbital and topiramate are taken at the same time or if the Epilim starting dose is high or has been suddenly increased.
- Changes in the amount of ammonia in the blood. Symptoms of this condition are being sick, problems with balance and co-ordination, feeling lethargic or less alert.
- Feeling shaky (tremor), sleepy or unsteady when walking or jerky muscle movements.
- Feeling tired or confused with loss of consciousness sometimes accompanied by hallucinations or fits.
- Blisters with the skin flaking away.
- Rapid, uncontrollable movement of the eyes.
- An increase in the number and severity of convulsions.

**Tell your doctor or pharmacist if any of the following side effects get serious or lasts longer than a few days, or if you notice any side effects not listed in this leaflet:**

- Feeling sick (nausea), being sick (vomiting), stomach ache or diarrhoea, especially when starting treatment. This may be helped by taking the tablets with food.
- Swelling of gums or sore mouth
- Fainting
- Hearing loss

- Nail and nail bed disorders
- Skin problems such as rashes. These happen rarely, but more often in people also taking lamotrigine.
- Acne
- Hair loss which is usually temporary. When it grows back it may be more curly than before.
- Hair disorders (changes in texture, colour or growth)
- Increased levels of some hormones (androgens), which may lead to increased hair growth on the face, breasts or chest, acne or thinning hair.
- Skin rash caused by narrow or blocked blood vessels (vasculitis)
- Changes in women's periods and increased hair growth in women
- Breast enlargement in men
- Swelling of the feet and legs (oedema)
- Weight gain – as your appetite may be increased
- Kidney disease
- Kidney problems, bedwetting or increased need to pass urine
- Blood in the urine
  - Headache
  - Seeing or hearing things that are not there (hallucinations)
  - Aggression, agitation, disturbance in attention, abnormal behaviour, restlessness/hyperactivity, and learning disorder
  - Tingling or numbness in the hands and feet
  - Lowering of normal body temperature
  - Abnormal blood clotting factors
  - Muscle pain and weakness (rhabdomyolysis)
  - Obesity

### **Bone disorders**

There have been reports of bone disorders including osteopenia and osteoporosis (thinning of the bone) and fractures. Check with your doctor or pharmacist if you are on long-term anti-epileptic medication, have a history of osteoporosis, or take steroids.

### **Blood tests**

Epilim can change levels of liver enzymes, salts or sugars shown up on blood and urine tests.

### **Male fertility**

Taking Epilim can be a contributing factor in male infertility.

### **Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

#### United Kingdom

You can also report side effects directly via the Yellow Card Scheme at:

[www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store.

#### Malta

You can also report side effects directly via ADR Reporting:

[www.medicinesauthority.gov.mt/adrportal](http://www.medicinesauthority.gov.mt/adrportal)

By reporting side effects you can help provide more information on the safety of this medicine.



## 5. How to store Epilim

Keep out of the sight and reach of children.

Do not take this medicine after the expiry date shown on the blister and carton after EXP. The expiry date refers to the last day of that month.

Do not remove the tablets from the foil until just before you take them. Do not cut the blister strips. Store in a dry place below 30°C.

Medicines should not be disposed of via household wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Epilim Gastro-resistant Tablets contain

- Each 200mg gastro-resistant tablet contains 200mg of the active substance, sodium valproate.
- Each 500mg gastro-resistant tablet contains 500mg of the active substance, sodium valproate.
- The other ingredients are povidone (E1201), talc, calcium silicate (E552), magnesium stearate (E572), hypromellose (E464), citric acid monohydrate (E330), macrogol 6000, polyvinyl acetate phthalate, diethyl phthalate, stearic acid (E570), titanium dioxide (E171), amaranth aluminium lake (E123), indigo carmine lake (E132) and hydroxypropyl cellulose (E463).

### What Epilim Gastro-resistant Tablets look like and contents of the pack

Epilim tablets are round and lilac coloured. The tablets are supplied in blister packs of 100.

### Marketing Authorisation Holder and Manufacturer

#### Marketing Authorisation Holder

Sanofi

One Onslow Street

Guildford

Surrey

GU1 4YS

UK

Tel: 0845 372 7101

email: uk-medicalinformation@sanofi.com

#### Manufacturer

Sanofi-aventis S.A,

Carretera C-35 (La Batlloria-Hostalric), Km 63.09

17404 Riells i Viabrea (Girona), Spain

This leaflet does not contain all the information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

**This leaflet was last revised in March 2018**

**© Sanofi, 2006-2018**

There are two organisations that will also be happy to try and answer any general questions on epilepsy. They can be contacted at:

**Epilepsy Action**, New Anstey House, Gate Way Drive, Yeadon, Leeds, LS19 7XY

Telephone: 0808 800 5050

Website: [www.epilepsy.org.uk](http://www.epilepsy.org.uk)

**National Society for Epilepsy (NSE)**, Chesham Lane, Chalfont St Peter, Bucks, SL9 0RJ

Telephone: 01494 601400

Website: [www.epilepsynse.org.uk](http://www.epilepsynse.org.uk)