This area must be identical on both sides!

- if you suffer from migraine, asthma, or kidney problems
- ▶ if you suffer from epilepsy (see "Other medicines and Primolut N")
- if you have an inflammation of your veins (superficial phlebitis)
- need to check your blood pressure and make sure you are not pregnant. You may also need additional checks, such as a breast examination, that will be specific to your medical ▶ if you have varicose veins

2. What you need to know before you take Primolut N

▶ if you are pregnant or if you think you might be pregnant

▶ if you have ever suffered migraine with visual disturbance

needs and/or concerns

▶ if you are breast-feeding

liver is not vet working normally

▶ if you have (or have ever had) liver tumours

sex hormones (including breast cancer)

if you have diabetes with damaged blood vessels

vellowing of the skin (idiopathic jaundice of pregnancy)

Talk to your doctor or pharmacist before taking Primolut N

mbosis, a heart attack or a stroke at a voung ag

itching of the whole body (nruritus of nregnancy)

or:

were pregnant:

if you smoke

Warnings and precautions

during treatment

If you are overweight (BMI ≥ 30 kg/m²)

if you have high blood pressure

Do not take Primolut N:

Your doctor will discuss your medical and family history with you. Your doctor will also

if you are allergic to norethisterone or any of the other ingredients of this medicine (listed in section 6)

▶ if you have ever had a problem with your blood circulation. This includes a blood clot

if you have any symptoms of a blood clot, such as chest pain, unexplained and often sudden shortness of breath and/or cough

▶ if you have any condition which makes you more at risk of a blood clot (thrombosis)

▶ if you have (or are recovering from) a liver disease and the blood tests show that your

if you have any type of cancer which might be made worse by exposure to female

▶ if you have a condition called endometrial hyperplasia which has not been treated.

In addition, do not take Primolut N if you have had any of the following conditions when you

if you have diabetes. Primolut N can produce changes in blood sugar levels. If you are diabetic, your doctor will check your blood sugar before starting treatment and regularly

if you have a heart valve disorder or a certain heart rhythm disorder (heart problems)

• if you have had a thrombosis/embolism or anyone in your close family has had a

• if you have problems with genital bleeding for which the cause is not yet known

→ Tell your doctor if any of these apply to you and do not take Primolut N.

(thrombosis) in the legs (deep ven thrombosis), lungs (pulmonary embolism), heart (heart attack), brain (stroke) or any other parts of the body

- if anyone in your immediate family has had breast cancer
- if you have previously had a condition called chloasma where the skin on your face may develop brownish blotches. You may be advised to avoid exposure to the sun and to ultraviolet light while you are taking Primolut N
- ▶ if you have previously suffered from depression
- ▶ if you or someone in your close family has ever had high blood levels of cholesterol or triglycerides (fatty substances)
- If you have a disease of the liver or gall bladder
- if you have certain rare medical conditions such as systemic lupus erythematosus (SLE), sickle cell disease, Crohn's disease or ulcerative colitie if you have haemolytic uremic syndrome (HUS)
- ▶ if you have a condition that occurred for the first time or worsened during pregnancy or previous use of sex hormones (e.g. hearing loss, porphyria, or Sydenham's chorea) if you have hereditary angioedema. Consult your doctor immediately if you experience
- symptoms of angioedema such as swollen face, tongue or throat, and/or difficulty swallowing, or hives, together with difficulty breathing. Products containing oestrogens may induce or worsen symptoms of angioedema

if you have an intolerance to some types of sugar (galactose intolerance, Lapp) lactase deficiency or glucose-galactose malabs

If you are using other medications as mentioned in "Other Medicines and Primolut N".

→ Tell your doctor before you take Primolut N if any of these applies to you. Also tell your doctor if any of these conditions develop or worsen while you are taking Primolut N, because you may need to stop taking it.

Primolut N and blood clots:

The main ingredient in Primolut N (progestogen) is partly converted into gestrogen so you should also consider the general warnings given for combined oral contraceptive pills ("the Pill").

Do not take Primolut N if you have a **blood clot** or have any medical condition which makes you more at risk of developing clots.

The risk of blood clots occurring in the veins and arteries is slightly greater in women who The lists of blood clobs becaring in time Veits and arenes is sugnity greater in wonch wino take the combined oral contractive pill than in wonces strokes, heart attacks and blood lobs fully recover from such blood clots, which can cause strokes, heart attacks and bleding into the brain (subscription).

You are more at risk of having a blood clot: as you get older

- if you're off your feet for a long time because of major surgery, injury or illness.
- ▶ if you smoke

۲

- ▶ if you or any of your close family have had blood clots
- if you are overweight (BMI ≥ 30 kg/m²)

pain and tightness in the chest sudden weakness or numbness in one side or part of your body dizziness or fainting.

See a doctor as soon as possible if you notice any possible signs of blood clot. Do not take any more Primolut N until your doctor says you can.

if you have a heart valve disorder or a particular type of irregular heartbeat (atrial

if you have certain medical conditions such as systemic lupus erythematosus (SLE), sickle cell disease, Crohn's disease or ulcerative colitis

if you are going to be immobilised for a long time (e.g. if you need bed-rest after an
accident or operation, or if you have a plaster cast on a broken leg)

any sudden changes to your evesight (such as loss of vision or blurred vision)

any sudden changes to your hearing, speech, sense of smell, taste or touch

To reduce the risk of blood clots, treatment with Primolut N must be stopped

• a migraine for the first time or one that is worse than normal

→ Tell your doctor if any of these apply to you. Taking Primolut N may add to this risk so

Primolut N and cancer

if you have a disorder of blood fat (lipid) metabolism

if you have a blood disorder

if you suffer from migraines

fibrillation)

if you have high blood pressure

▶ if you have recently had a baby ▶ if you have diabetes

it may not be suitable for y

before any surgery to the legs

Signs of a blood clot include:

pain or swelling in your leg

breathlessness

stabbing pain when you breather

couching for no apparent reason

six weeks before any planned major operation

before medical treatment for varicose veins

unusually frequent or severe headaches

If you have **breast cancer,** or have had it in the past, you should not take combined oral I you must be the contract of mark indices that the pass, you should not use contained to the contraceptives (the Pill). The Pill slightly increases your risk of breast cancer. This risk goes up the longer you're on it, but returns to normal within about 10 years of stopping it. Because breast cancer is rare in women under the age of 40, the extra cases of breast cancer in current and recent Pill users is small. For example:

Of 10.000 women who have never taken the Pill. about 16 will have breast cancer by the time they are 35 years old.

→ Please turn over

7/21/2017 11:23:06 AM

hnology Berlin sggwx material-no.: 85996428 olut N 5MG TAB GB

Peter Wittmuetz

85996428 01.indd 1

Package leaflet: Information for the user Primolut[®] N 5 mg tablets (norethisterone)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for

Keep this leaflet. You may need to read it again

- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours. If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Primolut N is and what it is used for 2. What you need to know before you take Primolut N

How to take Primolut N

۲

4. Possible side effects 5. How to store Primolut N

6. Contents of the pack and other information

- 1. What Primolut N is and what it is used for
- Primolut N contains norethisterone, which belongs to a group of medicines called progestogens, which are female hormones.
- Primolut N can be used in several different circumstances:
- to treat irregular, painful or heavy periods
- to treat endometriosis (where tissue from the lining of the womb is present in places) where it is not normally found)
- ▶ to treat premenstrual syndrome (also known as premenstrual tension, PMS or PMT)
- to delay periods



5MG TAB GB

the time they are 45 years old.

Your risk of breast cancer is higher if:

you are overweight (BMI ≥ 30 kg/m²)

liver tumours, but this is rare.

treat epilepsy.

Other medicines and Primolut N

Sedative drugs called barbiturates.

St lohn's wort to treat depression.

Other things you should know:

Pregnancy and breast-feeding

Driving and using machines

medicine.

The antibiotic drugs rifampicin or rifabutin

called ciclosporin which is often used after an organ transplant

Primolut N is unlikely to affect your ability to drive or use machines.

An antifungal drug called ariseofulvin.

have breast cancer by the time they are 35 years old.

have breast cancer by the time they are 45 years old.

• Of 10,000 women who take the Pill for 5 years in their early thirties, about 110 will

> you have a close relative (mother, sister or grandmother) who has had breast cancer

→ See a doctor as soon as possible if you notice any changes in your breasts, such as

Very rarely, the Pill has been linked with some forms of liver cancer in women who take it for a long time. These may lead to bleeding in the abdomen.

→ See a doctor as soon as possible if you get severe pain in your stomach that does not go away, or yellow skin or eyes (jaundice). You may need to stop taking Primolut N.

→ Tell your doctor if you are taking, have recently taken or might take any other medicines.

▶ Phenytoin, oxcarbazepine, primidone or carbamazepine. These drugs are often used to

Some other medicines may be affected by Primolut N. Tell your doctor if you are taking a drug

Taking Primolut N can affect the results of some blood and urine tests. Tell your doctor that you are taking Primolut N if you are asked to provide a blood or urine sample.

Once you have finished taking a course of Primolut N, you will usually have a menstrual

bleed (period) 2-3 days after taking your last tablet. If you do not have a period, you must make sure that you are not pregnant before taking any more tablets.

Do not take Primolut N if you are pregnant or breast-feeding. If you think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this

Some medicines may affect the way Primolut N works. Tell your doctor if you are taking:

Taking the Pill has also been linked to liver diseases, such as jaundice and non-cancerous

dimpling of the skin, changes in the nipple or any lumps you can see or fe

This area must be identical on both sides! 🍄

▶ Of 10,000 women who take the Pill for 5 years in their early twenties about 17–18 will Primolut N contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact Of 10.000 women who have never taken the Pill. about 100 will have breast cancer by your doctor before taking this medicinal product.

3. How to take Primolut N

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. The number of tablets that you need to take and the number of days per month when you

need to take them will depend on why the doctor has prescribed Primolut N. A common dosage would be 2-3 tablets each day. For some conditions Primolut N has to be taken every day, but this is not always the case.

Ask your doctor or pharmacist, if you are not sure about the number of tablets that you need to take, when they should be taken or how long you should take them for. Swallow the tablets whole with a drink of water.

If you take more Primolut N than you should

Taking too many tablets is unlikely to cause serious problems. If you take too many, contact your doctor who will tell you what to do.

If you forget to take Primolut N

If you forget a dose, wait until it is time to take the next prescribed dose. Do not take the missed dose. If you are worried, contact your doctor or pharmacist. If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Reasons for stopping Primolut N immediately:

Stop taking Primolut N and speak to your doctor immediately if you experience any of the following:

- migraine for the first time
- unusually bad headaches, occurring more often than before
- sudden changes to your eyesight, hearing or speech
- sudden changes to your senses of smell, taste or touch
- symptoms of blood clot formation or symptoms of inflammation of the veins combined
- with the formation of blood clots (thrombophlebitis)
- unusual pains in your leg(s)
- unusual swelling of your arms or legs
- > sharp pains in your chest or sudden shortness of breath
- > crushing pains or feelings of heaviness or tightness in your chest
- ▷ coughing for no apparent reason
- > one side of your body suddenly becoming very weak or numb

- Primolut N must also be stopped immediately if vou become pregnant
 - vou develop jaundice or other liver problems
 - you develop itching (pruritus)
 - your doctor finds that your blood pressure is too high

General side effects:

Side effects that have been reported with Primolut N are listed below according to the frequency with which they occur.

Very common side effects (These may affect more than 1 in 10 people)

Vaginal bleeding, including spotting.

- Periods that are much shorter than normal and where blood flow is reduced. Common side effects
- (These may affect up to 1 in 10 people)
- Headache
- Swelling

(These may affect up to 1 in 100 people)

(These may affect up to 1 in 1,000 people) Allergic reactions including skin rash or "hives"

Visual disturbances

- (Frequency not known: frequency cannot be estimated from the available data)
- Worsening of depression
- Dizziness Abdominal pain
- Jaundice
- Cholestasis

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine. United Kingdom

Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard

Malta ADR Reporting

Website: www.medicinesauthority.gov.mt/admortal

5. How to store Primolut N

Keep this medicine out of the sight and reach of children. Store in the original carton.

Do not use this medicine after the expiry date which is stated on both the outer carton and on each blister strip of tablets after EXP. The expiry date refers to the last day of that month Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Primolut N contains

Each tablet contains 5 mg of the active ingredient, norethisterone. The other ingredients are lactose, maize starch and magnesium stearate (E572).

What Primolut N looks like and contents of the pack

Each pack contains 30 tablets (3 foil blister packs containing 10 tablets each). Each white tablet has 'AN' embossed in a regular hexagon on one side

Marketing Authorisation Holder and Manufacturer Marketing authorisation holder:

Bayer plc 400 South Oak Way Reading RG2 6AD

United Kingdom Manufacturers:

Baver AG. Mullerstrasse 178, 13353 Berlin, Germany

Bayer Weimar GmbH & Co. KG, 99427 Weimar, Germany This leaflet was last revised in August 2017. Product licence number:

PL 00010/0553

Baver 85996628

۲

۲

- Feeling sick (nausea)
- Absence of a period

Uncommon side effects

Migraine

Rare side effects

Very rare side effects

Difficulty in breathing

Other side effects

(These may affect up to 1 in 10,000 people)

1.3.2