



Package leaflet: Information for the user

Scheriproct® Ointment

(1.9 mg prednisolone hexanoate / 5.0 mg cinchocaine hydrochloride per 1 g)

Scheriproct® Suppositories

(1.3 mg prednisolone hexanoate / 1.0 mg cinchocaine hydrochloride per suppository)

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Scheriproct is and what it is used for

2. What you need to know before you use Scheriproct

3. How to use Scheriproct

4. Possible side effects

5. How to store Scheriproct

6. Contents of the pack and other information

1. What Scheriproct is and what it is used for

Scheriproct contains a substance which reduces inflammation (prednisolone) and a local anaesthetic (cinchocaine) which relieves pain.

This medicine is used for the relief of the inflammation, swelling, itching and soreness of piles (haemorrhoids) and to relieve itching of the anus (back passage). It is used short-term usually for 5 to 7 days.

2. What you need to know before you use Scheriproct

Do not use Scheriproct:

- if you are allergic to prednisolone hexanoate, cinchocaine hydrochloride, other local anaesthetics or any of the other ingredients of this medicine (listed in section 6).
- if you have a viral infection (e.g. herpes, shingles, chicken-pox)
- if you have any bacterial or fungal infections of the skin or elsewhere for which you are not receiving treatment.

Warnings and precautions

Talk to your doctor or pharmacist before using Scheriproct

- Long-term continuous treatment should be avoided because it increases the possibility of side effects. This is particularly important for infants and small children because continuous treatment with Scheriproct for long periods may reduce the activity of the adrenal glands and so lower resistance to disease. Also, long-term treatment can cause the skin to thin and deteriorate in the affected area (see section 4) and some of the medicine may be absorbed into the blood stream.

- If the area treated with Scheriproct is also infected your doctor should prescribe another medicine, to use with Scheriproct, to treat the infection.
- Do not use a waterproof dressing to cover areas where you have applied the cream.

Other medicines and Scheriproct

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicines may increase the effects of Scheriproct and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. There may be a very small risk to the development of a baby in pregnant women treated with Scheriproct. As with most medicines, this risk is likely to be greatest during the first 3 months of pregnancy.

Driving and using machines

Scheriproct has no influence on the ability to drive and use machines.

3. How to use Scheriproct

Always use this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. Do not use Scheriproct for more than 7 days.

Always wash your hands before and after applying Scheriproct.

How to use Scheriproct Ointment:

- Before you use the ointment, gently but thoroughly wash and dry the anus (back passage) and the skin around it.
- If the ointment is to be used for the area **around** the anus:
 - Squeeze a small quantity (about the size of a pea) on to the top of your finger.
 - Spread it gently over the skin and just inside the anus.
 - Do not rub it in.
- If the ointment is to be used **inside** the anus:
 - Put the separate nozzle (applicator) onto the tube.
 - Squeeze the tube until the nozzle is full of ointment. The amount of ointment in the nozzle is one dose.
 - Insert the nozzle very carefully into the anus until the whole length of the nozzle is inside.
 - Then, while squeezing the tube gently, withdraw the nozzle.
 - Wash the nozzle carefully in hot soapy water and rinse thoroughly.

Generally, the ointment should be applied twice a day, but it may be applied three or four times on the first day, to obtain quick relief.

438850/GB/9



Bayer HealthCare Manufacturing S.r.l. (Segrate)

Artwork date: 18/04/2017
Version n°: 03
Page: 1/2

Technical drawing:	Item n°:	Replaces:
200 x 305 mm	438850/GB/9	438850/GB/8
Laetus-code n°:	Client:	
Bar-code n°:	599	
Name:	Country:	
Scheriproct supp. / ointment	Great Britain	
Colors:		
Black		Recto - Font size 9 pt
Approved:		

cod 135913 sdp FRGBRBBB | tel 02 38 1061

1. Before you insert a suppository find the small tear in the foil packet and remove the covering foil, by tearing it in half.
2. If the suppositories have become softened, owing to warm temperature, they can be hardened by putting them into cold water before you remove the covering foil.
3. Insert the whole suppository into the anus.
4. To make insertion easier, either stand with one foot raised on a chair or squat down.

If you use more Scheriproct than you should or if you accidentally swallow some Scheriproct

If you forget to use Scheriproct

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

Allergic skin reactions may occur in rare cases. Castor oil, one of the ingredients of Scheriproct Ointment may cause a skin reaction.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

- ▶ Do not store Scheriproct Ointment above 25°C.
- ▶ Store Scheriproct Suppositories in a refrigerator (2°C to 8°C).
- ▶ Do not use Scheriproct after the expiry date which is stated on the pack after EXP. The expiry date refers to the last day of that month.
- ▶ Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.


► The active substances are prednisolone hexanoate and cinchocaine hydrochloride. 1 g of Scheriproct Ointment contains 1.9 mg prednisolone hexanoate and 5.0 mg cinchocaine hydrochloride

- **Scheriproct Suppositories** are yellowish-white and are supplied in packs of 12 suppositories.

The suppositories are manufactured by:
Istituto de Angeli S.r.l., Reggello, Italy
or Bayer HealthCare Manufacturing S.r.l.
Via E. Schering 21
20090 Segrate (Milan)
Italy

Bayer

438850/GB/9

		Bayer HealthCare Manufacturing S.r.l. (Segrate)		Artwork date: 18/04/2017 Version n°: 03 Page: 2/2	
Technical drawing): 200 x 305 mm		Item n°: 438850/GB/9		Replaces: 438850/GB/8	
Laetus-code n°: Bar-code n°: 816		Client: 599			
Name: Scheriproct supp. / ointment		Country: Great Britain			
Colors: Black		Verso - Font size 9 pt			
Approved:					

cod 135913 **gdp** PREPRESS | tel. 02.38.106.1