

Diazepam 2 mg, 5 mg and 10 mg Tablets

Diazepam

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effect not listed in this leaflet. See section 4.

What is in this leaflet

1. What Diazepam Tablets are and what they are used for
2. What you need to know before you take Diazepam Tablets
3. How to take Diazepam Tablets
4. Possible side effects
5. How to store Diazepam Tablets
6. Contents of the pack and other information

1. What Diazepam Tablets are and what they are used for

The name of your medicine is Diazepam 2 mg, 5 mg or 10 mg Tablets (referred to as Diazepam Tablets in this leaflet). Diazepam Tablets contain the active ingredient diazepam, which belongs to a group of medicines called benzodiazepines. Diazepam Tablets help in the treatment of anxiety, muscle spasms and convulsions (fits).

Diazepam Tablets are used to treat a number of conditions, including:

In adults

- short term relief (2-4 weeks only) of severe anxiety, which is an emotional state where you may sweat, tremble, feel anxious and have a fast heart beat and may occur alone or with insomnia (trouble sleeping) or mental health problems
- helping muscles relax and for muscle spasm and cerebral palsy (a condition affecting the brain which causes movement problems and rigidity or stiffness)
- epilepsy (when taken with other medicines)
- patients with the symptoms of alcohol withdrawal
- helping to relax nervous dental patients.

In children

- helping to treat tension and irritability caused by cerebral spasticity (a condition associated with a disease or trauma affecting the brain or spinal cord which causes weakness, un-coordinated movements, rigidity and stiffness)
- helping to treat muscle spasm caused by tetanus (when taken with other medicines).

Both adults and children can take Diazepam Tablets before an operation to help with relaxation and to cause sleepiness.

2. What you need to know before you take Diazepam Tablets

Do not take Diazepam Tablets

- if you are allergic to diazepam or any of the other ingredients of this medicine (listed in section 6).
- if you have a phobia (a fear of a particular object or situation) or other mental illness
- if you have breathing problems, which may be severe, including slow and/or shallow breathing
- if you have myasthenia gravis (a condition which causes muscles to weaken and tire easily)
- if you suffer from sleep apnoea (a sleep disorder where you have abnormal pauses in breathing during sleep)
- if you have severe liver disorders
- if you have porphyria (an inherited condition causing skin blisters, abdominal pain and brain or nervous system disorders)
- if you suffer from depression (with or without anxiety) or hyperactivity
- if you are planning a pregnancy or are pregnant (see below *Pregnancy and breast-feeding*).

Warnings and precautions

Talk to your doctor or pharmacist before taking Diazepam Tablets

- if you have long term kidney, liver, heart or respiratory problems
- if you have a personality disorder
- if you have a history of alcoholism or drug abuse
- you have a history of seizures (fits)
- if someone close to you has recently died
- if you have low blood levels of a protein called albumin
- if you are elderly (risk of confusion or clumsiness, causing fall or injury)
- if you have a poor blood supply to the brain (arteriosclerosis)
- If you have breathing difficulties
- If you smoke
- If you suffer from depression
- If you have suicidal thoughts
- If you have epilepsy or a history of seizures

Other considerations

- **Mental side effects** – contact your doctor if you experience side effects such as agitation, hyperactivity, restlessness, aggressiveness, nightmares or hallucinations. These side effects are more likely to occur in children or the elderly.
- **Amnesia** (total or partial memory loss) – you could experience amnesia when taking this medicine. To reduce the risk you should ensure that you will be able to have uninterrupted sleep of 7-8 hours. Amnesia is more likely to occur when taking high doses of diazepam.
- **Dependence** - when taking this medicine there is a risk of dependence, which increases with the dose and duration of treatment and also in patients with a history of alcoholism and drug abuse. Therefore, you should take Diazepam Tablets for a short a period of time as possible.
- **Tolerance** - if after a few weeks you notice that the tablets are not working as well as they did when first starting treatment, you should speak to your doctor.

- **Withdrawal** - treatment should be gradually withdrawn. Withdrawal symptoms occur with Diazepam Tablets even when normal doses are given for short periods of time. See Section 3, 'If you stop taking Diazepam Tablets'.

Other medicines and Diazepam Tablets

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines, including medicines obtained without prescription.

This is especially important if you are taking the following:

- **antidepressants** (e.g. fluvoxamine, fluoxetine)
- **antipsychotics** such as clozapine (to treat mental problems)
- **antihistamines** (to treat allergies)
- **general anaesthetics**
- **sedatives** (used to give calming effects)
- **hypnotics** (to help you sleep)
- **erythromycin** (an antibiotic)
- **muscle relaxants** (e.g. suxamethonium, tubocurarin)
- some **strong pain killers** such as **morphine** (opioids) may give you a heightened sense of well being when taken with diazepam, which can increase your desire to continue taking these medicines (dependency) or can make you very sleepy.
- barbiturates such as **phenobarbital** (to treat epilepsy and mental disorders)
- **medicines to lower high blood pressure, diuretics** (water tablets), **nitrates** (for heart conditions) as these could lower your blood pressure too much.
- **antacids** (reduces stomach acid) may slow down absorption of diazepam in the body.

Taking these medicines with diazepam could affect your mental status, make you very sleepy and suppress your breathing and blood pressure.

- **disulfiram** (to treat alcohol addiction). Taking this medicine with diazepam could make you very sleepy and can cause diazepam to be removed from the body more slowly than usual.
- **medicines for epilepsy** e.g. phenobarbital, phenytoin and carbamazepine, sodium valproate, (diazepam can affect the blood levels of these medicines). Diazepam can furthermore affect how phenytoin works.
- **theophylline** (to treat asthma and other breathing disorders), as it can weaken the effect of diazepam. As this can cause diazepam to be removed from the body more quickly than usual.
- **cimetidine, omeprazole or esomeprazole** (stomach acid reducing medicines), as these can cause diazepam to be removed from the body more slowly than usual.
- **rifampicin**, to treat infections (an antibiotic) as this can cause diazepam to be removed from the body more quickly than usual. The effect of diazepam can be weakened.
- **amrenavir, atazanavir, ritonavir, delavirdine, efavirenz, indinavir, nelfinavir or saquinavir** (antivirals), **fluconazole, itraconazole, ketoconazole or voriconazole** (anti-fungal medicines) as these can cause diazepam to be removed from the body more slowly than usual and therefore increase the risk of side effects. As these can make you feel sleepy for longer or cause difficulty breathing.
- **isoniazid** (used to treat tuberculosis), as it can cause diazepam to be removed from the body more slowly than usual.
- **oral contraceptives**, as they can slow down the removal of diazepam from the body and increase its effect. Breakthrough bleeding can occur when taking diazepam and oral contraceptives together, but the contraceptive protection is not reduced.
- **cisapride** (used to treat stomach problems), as it can cause diazepam to be removed from the body more slowly than usual.
- **corticosteroids** (medicines used to treat inflammation in the body) as they can weaken the effect of diazepam.
- **levodopa** (used to treat Parkinson's disease). Diazepam can reduce the effect of levodopa.
- **valproic acid** (used to treat epilepsy and mental disorders) as it can slow down the removal of diazepam from the body and increase its effect.
- **ketamine** (an anaesthetic) as diazepam increases the effect of ketamine.
- **lofexidine** (to help relieve symptoms when you stop taking opioids)
- **nabilone** (to treat nausea and vomiting)
- **alpha blockers or moxonidine** (to lower high blood pressure)

Diazepam Tablets and food and drink

Do not drink alcohol while you are taking Diazepam Tablets. Alcohol may increase the sedative effects of Diazepam Tablets and make you very sleepy.

Grapefruit juice may increase the amount of diazepam in your blood. If you are elderly, suffer from cirrhosis or any of the conditions listed in section 2, this could possibly increase the sedative effects of Diazepam Tablets and you should speak to your doctor or pharmacist.

Drinks containing caffeine may reduce the effects of diazepam.

Other special warnings

- If you see another doctor or visit a hospital, remember to tell them what medicines you are already taking.
- If you are going to have an operation please tell the anaesthetist that you are taking Diazepam Tablets as this may affect some of the drugs he/she may use.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.

You should not take Diazepam Tablets if you are pregnant, planning to become pregnant or are breast-feeding. If you take Diazepam Tablets late in your pregnancy or during labour, your baby might have a low body temperature, floppiness and breathing difficulties.

If taken regularly during late pregnancy, your baby may develop withdrawal symptoms.

Diazepam can pass into breast milk. Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Diazepam Tablets can make you sleepy, forgetful, have poor co-ordination along with other side effects that can affect everyday activities (see *Possible side effects*). You should not drive, operate machinery or take part in such activities where, if affected, you could put yourself or others at risk.

The medicine can affect your ability to drive as it may make you sleepy or dizzy.

- Do not drive while taking this medicine until you know how it affects you.
- It is an offence to drive if this medicine affects your ability to drive.
- However, you would not be committing an offence if:
 - the medicine has been prescribed to treat a medical or dental problem, and;
 - you have taken it according to the instructions given by the prescriber or in the information provided with the medicine, and;
 - it was not affecting your ability to drive safely.

Talk to your doctor or pharmacist if you are not sure whether it is safe for you to drive while taking this medicine.

Diazepam Tablets contain lactose

Diazepam Tablets contain lactose (a type of sugar). If you have been told that you have an intolerance to some sugars contact your doctor before taking this medicine.

3. How to take Diazepam Tablets

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You should not take Diazepam Tablets for longer than 4 weeks.

The tablets should be swallowed whole with a glass of water.

Recommended dose

Adults

- **Anxiety or mental health problems:** 5mg-30mg each day, in divided doses.
- To help you sleep: 5mg-15mg at bedtime.
- To help cerebral palsy or other spasticities: 5mg-60mg each day, in divided doses.
- To help control muscle spasm: 5mg-15mg each day, in divided doses.
- To help epilepsy: 2mg-60mg each day, in divided doses.
- To help with alcohol withdrawal symptoms: 5mg-20mg, which may be repeated after 2 to 4 hours if necessary.
- Before dental treatment: 5mg the night before treatment, 5mg on waking and 5mg two hours before the appointment.
- Before an operation: 5mg-20mg

Elderly or frail

If you are elderly or frail you are likely to be more sensitive to the effects of Diazepam Tablets, such as confusion, and your doctor will give you much lower doses. The dose should not be more than half the adult dose.

Renal and/or hepatic impairment

If you have liver or kidney problems you may also be given a lower dose.

Use in children

For tension and irritability in cerebral spasticity: 5mg-40mg each day, in divided doses.

If your doctor has given your child Diazepam Tablets to take before an operation, the usual dose is 2mg-10mg.

If you take more Diazepam Tablets than you should

If you (or someone else) swallow a lot of tablets at the same time, or you think a child may have swallowed any, contact your nearest hospital casualty department or tell your doctor immediately. Signs of an overdose include clumsiness and loss of coordination, feeling sleepy or deep sleep, speech problems, irregular or slow heartbeat, uncontrolled eye movement, muscle weakness or excitement. An extreme overdose may lead to coma (unrousable unconsciousness), reflex problems and breathing difficulties.

If you forget to take Diazepam Tablets

If you forget to take a dose, just take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

If you stop taking Diazepam Tablets

- Do not stop taking your medicine without telling your doctor as she/he may wish to gradually reduce the number of tablets you take before stopping them completely. If you stop taking Diazepam Tablets suddenly, you may experience unpleasant side effects including depression, nervousness, irritability, sweating or diarrhoea. If you have been taking a high dose, you may occasionally experience confusion, convulsions or unusual behaviour.
- Treatment should be gradually withdrawn otherwise the symptoms you are being treated for may return more intensely than before (rebound insomnia and anxiety). The risk of this happening is greater when you stop taking Diazepam Tablets suddenly. You may also experience mood changes, anxiety, restlessness or changes in sleep patterns.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines this medicine can cause side effects in some people, although not everybody gets them.

Serious side effects

If you notice any of the following, stop taking Diazepam Tablets and contact your doctor immediately:

Uncommon: may affect up to 1 in 100 people

- Respiratory depression (very slow and/or shallow breathing)

Rare: may affect up to 1 in 1,000 people

- Respiratory arrest (cessation of breathing)
- Unconsciousness
- Jaundice (yellowing of your skin or the white of your eyes)

Very rare: may affect up to 1 in 10,000 people

- Anaphylaxis (severe allergic reaction) with symptoms such as sudden wheezing, swelling of your lips, tongue and throat or body, rash, fainting or difficulties to swallow

Other side effects

Very common: may affect more than 1 in 10 people

- Drowsiness

Common: may affect up to 1 in 10 people

- Fatigue
- Withdrawal symptoms (for possible symptoms please see 'If you stop taking Diazepam Tablets' in Section 3)
- Confusion
- Loss of coordination of muscle movements (ataxia) and other movement disorders, tremor

Uncommon: may affect up to 1 in 100 people

- Muscle weakness
- Memory loss
- Difficulty in concentrating
- Balance disorders
- Dizziness
- Headache
- Slurred speech
- Stomach and intestinal problems such as nausea, vomiting, constipation, diarrhoea
- Increased salivation
- Allergic skin reactions in the form of itching, skin redness and swelling and skin rash.

Rare: may affect up to 1 in 1,000 people

- Mental side effects such as excitation, agitation, restlessness, irritability, aggressiveness, memory loss, delusion, rages, psychoses, nightmares or hallucinations. May be or become serious. These side effects are more likely to occur in children or the elderly. Talk to your doctor.
- Decreased alertness
- Depression
- Emotional withdrawal
- Insomnia (problems sleeping)
- Heart problems such as slow heartbeat (bradycardia), heart failure and cessation of heartbeat (cardiac arrest).
- Low blood pressure, fainting (syncope)
- Increased mucus in the lungs
- Dry mouth
- Increased appetite
- Changes in certain liver enzymes as seen in blood tests
- Lack of ability to urinate, loss of bladder control (leakage of urine)
- Breast enlargement in men
- Impotence, changes in sexual drive (libido)
- Blood disorders (you may develop sore throats, nose bleeds or infections)

Very rare: may affect up to 1 in 10,000 people

- Low levels of white blood cells (leukopenia)
- Higher level of a certain enzyme in the blood (transaminase)

Not known: frequency cannot be estimated from the available data

- Blurred vision, double vision and involuntary eye movements (these side effects disappear after you have stopped taking diazepam)

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard
By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Diazepam Tablets

Keep this medicine out of the sight and reach of children. Do not store above 25°C. Keep the container tightly closed and store in the original container or package in order to protect from light.

Do not use this medicine after the expiry date that is stated on the carton, bottle or blister after EXP. This date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Diazepam Tablets contain

Each tablet contains 2 mg, 5 mg or 10 mg of the active ingredient diazepam.

Diazepam 2 mg and 5 mg Tablets also contain colloidal silicon dioxide, maize starch, magnesium stearate and lactose. Diazepam 5 mg Tablets also contain yellow iron oxide (E172) and quinoline yellow lake (E104).

Diazepam 10 mg Tablets contain dextrin, maize starch, magnesium stearate, lactose and indigo carmine lake (E132).

What Diazepam Tablets look like and contents of the pack

Diazepam 2 mg Tablets: Flat white tablets with bevelled edges, 8 mm diameter. One face is embossed with "DZ" and "2" separated by a breakline. The reverse face is plain.

Diazepam 5 mg Tablets: Flat, pale yellow tablets with bevelled edges, 8 mm diameter. One face is embossed with "DZ" and "5" separated by a breakline. The reverse is plain.

Diazepam 10 mg Tablets: Flat, blue tablets with bevelled edges, 8 mm diameter. One face is embossed with "DZ" and "10" separated by a breakline. The reverse is plain.

Diazepam 2 mg, 5 mg and 10 mg Tablets are available in blister packs of 28 tablets and pots of 21, 100, 250, 500 and 1000 tablets.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Sovereign Medical, Sovereign House, Miles Gray Road, Basildon, Essex, SS14 3FR.

Manufacturer

Waymade Plc, Sovereign House, Miles Gray Road, Basildon, Essex, SS14 3FR.

This leaflet was last revised in July 2016