

9 mm

14 mm

Verso

11.5 mm

210 mm

11.5 mm

148 mm

3. HOW TO GIVE TIXYLIX TODDLER SYRUP

Always give Tixylix Toddler Syrup exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.
Using the measuring spoon provided, the following doses may be given 3 to 4 times daily, by mouth, as required. Allow 3 to 4 hours between doses.

Age	Dose
Under 3 months	Do not give Tixylix Toddler Syrup
3 to 12 months	One 5 ml spoonful
1 to 5 years	Two 5 ml spoonfuls

After use, wipe the neck of the bottle clean and make sure the safety cap is properly closed. You can help by keeping your child warm and giving plenty of fluids. Make sure the bedroom is well ventilated but not cold. If your child's condition does not get better in 3 days, talk to your doctor.

If you give more Tixylix Toddler Syrup than you should
Giving your child too much Tixylix Toddler Syrup may cause diarrhoea. If you have accidentally given your child more medicine than you should, or if you suspect that your child has taken extra medicine, please contact your doctor. Take along the bottle and the label, and any medicine that is left.

If you forget to give Tixylix Toddler Syrup
Do not give your child a double dose to make up for a forgotten dose.
If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Tixylix Toddler Syrup can cause side effects, although not everybody gets them. Tixylix Toddler Syrup may cause diarrhoea in overdose.

Reporting of side effects
If your child gets any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: www.mhra.gov.uk/yellowcard. By reporting side effects you can help provide more information on the safety of this medicine.

5. HOW TO STORE TIXYLIX TODDLER SYRUP

Keep out of the reach and sight of children.
Do not store above 25°C. Keep the container tightly closed.
Do not use Tixylix Toddler Syrup after the expiry date which is stated on the carton and the bottle label after <EXP>. The expiry date refers to the last day of that month.
Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

What Tixylix Toddler Syrup contains

- The active substance is glycerol (0.75 ml per 5ml dose).
- The other ingredients are liquid maltitol (Lycasin), hydroxyethylcellulose, sodium benzoate (E211), citric acid monohydrate, sodium citrate, natural flavours (vanilla, blackcurrant), purified water.

What Tixylix Toddler Syrup looks like and contents of the pack
Tixylix Toddler Syrup is a colour-free syrup.

The following pack sizes are available, not all pack sizes may be marketed:

- Amber-coloured glass bottle with child resistant cap, containing 100 ml or 150 ml. Measuring spoon provided.
- Amber-coloured polyethylene terephthalate bottle with child resistant cap, containing 150 ml. Measuring spoon provided.

Marketing Authorisation Holder
GlaxoSmithKline Consumer Healthcare (UK) Trading Limited, Brentford, TW8 9GS, U.K.

Manufacturer:
BCM Limited, Beeston, Nottinghamshire, NG90 2PR.

For any information about this medicinal product, please contact the Marketing Authorisation Holder.
This leaflet was last revised in January 2016

GB 316675 – XXX