



Artwork date: 16/05/2017
Version n°: 01
Page: 1/2

PZ (technical drawing):	2585A-4	Item n°:	REG	Replaces: 82092803
Laetus-code n°:	0013	Client:	044	
Bar-code n°:				
Name:	Ultraproct ointment	Country:	Great Britain (Meadow)	
Colors:	Black	Format label:		
Approved.:		Date:		



MEADOW

Ultraproct® Ointment
Ultraproct® Suppositories

Read all of this leaflet carefully before you start using this medicine.

- In this leaflet:**

1. What Ultraproct is and what it is used for
2. Before you use Ultraproct
3. How to use Ultraproct
4. Possible side effects
5. How to store Ultraproct
6. Further information

Ultraproct contains fluocortolone which belongs to a group of medicines known as the corticosteroids which reduce inflammation and itching. Ultraproct also contains a substance

Ultraproct can be used for the short-term (5-7 days) relief of inflammation, swelling, itching and soreness of piles (haemorrhoids) and to relieve itching of the anus (back passage).

This leaflet covers the following products:

- Ultraproct Suppositories
- Ultraproct Ointment

In this leaflet they are jointly referred to as Ultraproct.

Your doctor will prescribe the correct product to treat your specific condition.

Do not use Ultraproct if you:

- are **allergic** (hypersensitive) to local anaesthetics or any other ingredients of this medicine (see section 6)
- suffer from a **viral skin infection** such as cold sores, herpes, shingles or chickenpox
- have a **bacterial** or **fungal** infection for which you are not receiving treatment
- are **pregnant** or **breast-feeding**, unless recommended otherwise by your doctor

Tell your doctor if you believe any of the above conditions apply to you.

Take special care with Ultraproct:

Ultraproct **must not** come into contact with your eyes. Rinse with cold water if accidental contact occurs.

Avoid using Ultraproct for a long time because it increases the risk of side effects.

If the area treated with Ultraproct is also infected your doctor will prescribe another medicine to treat the infection.

Infants and Children

Ultraproct is not for use in infants and is not recommended for use in children under 12 years of age unless directed by a doctor.

Using dressings over the treatment area

Do not use waterproof dressings over the affected skin.

Taking other medicines

- Please tell your doctor if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- Some medicines may increase the effects of Ultraproct and your doctor may wish to monitor you carefully if you are taking these medicines (including some medicines for HIV: ritonavir, cobicistat).

Pregnancy and breast-feeding

Even at the usual dose, the possibility of a very small risk to the development of a baby in pregnant women treated with Ultraproct cannot be entirely ruled out. **Therefore, this medicine should not be used if you are pregnant, trying to become pregnant or are breast-feeding unless considered essential by your doctor.**

Driving and using machines

Ultraproct does not impair your ability to drive or operate machinery.

Important information about some of the ingredients of Ultraproct Ointment

Ultraproct Ointment contains castor oil and hydrogenated castor oil, which may cause skin reactions.

3. How to use Ultraproct

Wash your hands before and after using Ultraproct.

Do not use Ultraproct for more than 7 days.





PZ (technical drawing):	2585A-4	Item n°:	REG	Replaces: 82092803
Laetus-code n°:	0013	Client:	044	
Bar-code n°:				
Name:	Ultraproct ointment	Country:	Great Britain (Meadow)	
Colors:	Black	Format label:		
Approved.:		Date:		



If you or someone else accidentally swallows Ultraproct it is

Ointment: Do not store above 25° C.

REG

