

83,6 mm

**PACKAGE LEAFLET:
INFORMATION FOR THE USER**

Maalox® Plus Tablets

dried aluminium hydroxide gel 200mg
magnesium hydroxide 200mg
simeticone 25mg

sanofi aventis



**Is this leaflet hard to see or read?
Phone 01483 505515 for help**

Read all of this leaflet carefully because it contains important information for you

This medicine is available without prescription. However, you still need to use Maalox Plus Tablets carefully to get the best results from it.

- Keep this leaflet. You may need to read it again
- Ask your pharmacist if you need more information or advice
- You must contact a doctor if your symptoms worsen or do not improve
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist

In this leaflet:

1. What Maalox Plus Tablets are and what they are used for
2. Before you take Maalox Plus Tablets
3. How to take Maalox Plus Tablets
4. Possible side effects
5. How to store Maalox Plus Tablets
6. Further information

1. What Maalox Plus Tablets are and what they are used for



Maalox Plus Tablets contain three different medicines:

- The first two medicines are called dried aluminium hydroxide gel and magnesium hydroxide. They belong to a group of medicines called antacids.
- The other medicine is called simeticone. It belongs to a group of medicines called anti-foaming agents.

Maalox Plus Tablets is used for:

- The relief of indigestion (dyspepsia)
- Heartburn
- Wind (flatulence)

Maalox Plus Tablets work by lowering the amount of acid in your stomach. The anti-foaming agent in Maalox Plus Tablets helps to remove gas from your stomach.

2. Before you take Maalox Plus Tablets



Do not take Maalox Plus Tablets if:

- × You are allergic (hypersensitive) to dried aluminium hydroxide gel, magnesium hydroxide, simeticone or any of the other ingredients in Maalox Plus Tablets (see Section 6: Further information)
Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue
 - × You are feeling very weak and have no energy (debilitated)
 - × You have kidney problems
- Do not take this medicine if any of the above apply to you. If you are not sure, talk to your doctor or pharmacist before taking Maalox Plus Tablets.



Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes medicines you buy without a prescription, including herbal medicines.

Maalox Plus Tablets should not be taken at the same time as other medicines. This is because Maalox Plus Tablets can affect the way other medicines work. If Maalox Plus Tablets are taken within one hour of any other medicine, they can affect the uptake of that medicine into the blood.

Pregnancy and breast-feeding

Talk to your doctor or pharmacist before taking this medicine if:

- You are pregnant, might become pregnant, or think you may be pregnant
- You are breast-feeding or planning to breast-feed

Important information about some of the ingredients of Maalox Plus Tablets

Maalox Plus Tablets contain anhydrous glucose, sucrose and sorbitol (E420). This product may be harmful to teeth. If you have been told by your doctor that you cannot tolerate or digest some sugars, talk to your doctor before taking this medicine.

Turn
Over



3. How to take Maalox Plus Tablets

Taking this medicine

- Take this medicine by mouth
- Chew the tablets well before swallowing

Adults including the elderly

- Take one or two tablets 4 times each day
- Take 20 minutes to one hour after meals and at bedtime or as required

Children

Maalox Plus Tablets are not recommended for children
If the symptoms do not go away, talk to your doctor.

If you take more Maalox Plus Tablets than you should

Talk to your doctor or go to a hospital straight away. Remember to take any medicine that is left with you so the doctor knows what you have taken.

4. Possible side effects

Like all medicines, Maalox Plus Tablets can cause side effects, although not everybody gets them.

Tell your doctor or pharmacist if any of the following side effects get serious or lasts longer than a few days. Also tell them if you notice any side effects not listed in this leaflet:

- Constipation
- Diarrhoea

These side effects can happen, but are not common.

5. How to store Maalox Plus Tablets

Keep this medicine in a safe place where children cannot see or reach it

Do not use Maalox Plus Tablets after the expiry date which is stated on the carton and foil after EXP. The expiry date refers to the last day of that month.

Store below 30°C

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

What Maalox Plus Tablets contain

- Each tablet contains: 200mg of dried aluminium hydroxide gel, 200mg of magnesium hydroxide and 25 mg of simeticone
- The other ingredients are: swiss cream flavour, lemon flavour, magnesium stearate, talc, anhydrous citric acid, anhydrous glucose, saccharin sodium, sorbitol (E420), sorbitol liquid non-crystallising (E420), iron oxide yellow (E172), pregelatinised starch (maize starch), maize starch, sucrose and mannitol (E421)

What Maalox Plus Tablets looks like and contents of the pack

Maalox® Plus Tablets are white and yellow with “Maalox” embossed on one side. They taste of lemon/cream.

Maalox Plus Tablets are made in blister packs containing 10, 12, 20, 24, 30, 36, 40, 48, 50, 60, 70, 72, 80, 84, 90, 96, and 100 tablets or tubes containing 12, 24, 36, 48, 72 and 144 tablets. Not all pack sizes may be available.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Sanofi-aventis,
One Onslow Street,
Guildford,
Surrey, GU1 4YS, UK

Tel: 01483 505515

Fax: 01483 535432

email: uk-medicalinformation@sanofi-aventis.com

Manufacturer

Sanofi-aventis S.p.A.
Viale Europa, 11 - Origgio (VA), Italy

This leaflet does not contain all the information about your medicine. If you have any questions or are not sure about anything, ask your doctor or pharmacist.

This leaflet was last revised in 12/2010.

© sanofi-aventis 1989-2010